

The Effectiveness of Treatment for Adult Sex Offenders

Stephen Brake, Ph.D.¹

July 2010

Introduction

Evaluating the effectiveness of treatment for adult sex offenders is a difficult task. Designing research studies with adequate control groups is challenging and executing them is even more so. The fact that sex crimes are grossly under-reported lowers the (apparent) base rate of sex offending and limits conclusions that can be drawn about the significance of treatment effects. In addition, individual research studies require years to complete and so data about treatment effectiveness are very slow in coming. Meta-analyses of studies already completed are complicated by difficult decisions that must be made about what studies to include and how to compare treatment effectiveness among the disparate individual studies.

Because of these difficulties, conclusions about the effectiveness of treatment for adult sex offenders are difficult to reach. Yet it is important to know what the research tells us.

I examined 37 research studies concerning treatment outcome (29 individual studies and 8 meta-analyses) published over the last 26 years. The studies were easily accessible in peer-reviewed journals (or in some cases as government reports on websites) and required no particular expertise to interpret. I relied on the authors' conclusions about whether a finding was significant. I have also included here critiques of some of the research studies which were published by various authors in peer-reviewed journals. The survey is not intended to be a review of all articles ever written about the effectiveness of treatment for adult sex offenders but I have included all I could locate with reasonable effort and excluded none I found (other than a small few which clearly lacked control groups).

The form of treatment most commonly offered sex offenders in these studies is cognitive/behavioral or relapse prevention therapy (although other forms of treatment were sometimes the focus of the studies). The duration of treatment provided to offenders in the studies surveyed varies from just a few weeks to 4 years although 1-2 years was most common. The amount of time an offender was at risk following treatment ranged from 2 years to 28 years. (In some studies, the duration of treatment and amount of time at risk were difficult to determine.)

¹ Stephen C. Brake, Brake and Associates, 1181 South Parker Road, #103, Denver, CO, 80231, stephenbrake@stephenbrakeassociates.com.

Few definitive conclusions can be drawn from individual studies about the effectiveness of treatment. About half demonstrate that treatment can be effective in reducing recidivism and about half indicate that treatment had no statistically significant effect in reducing recidivism.

Table 1 summarizes the findings of the individual studies surveyed.




Table 1

Of the 29 individual studies surveyed, 12 (41%) concluded that some form of treatment significantly reduces recidivism (*Aytes, Olsen, Zakrajsek, Murray, and Ireson, 2001; Duve and Goldman, 2009; Hildebrand, deRuiter, and deVogel, 2004; Looman, Abracen, and Nicholaichuk, 2000; Maletzky, Tolan, and McFarland, 2006; Marshall and Barbaree, 1988; Marshall, Eccles and Barbaree, 1991; McGrath, Cumming, Livingston, and Hoke, 2003; McGrath, Hoke and Vojtisek, 1998; Nicholaichuk, Gordon, Gu, and Wong, 2000; West, Hromas, Wengler, and Suthers, 2000 - data from "Alaska"; Wilson, Picheca and Prinzgo, 2005*).

11 of the 24 studies (37%) concluded that treatment did not statistically reduce recidivism (*Barnoski, 2006; Davidson, 1984; Friendship, Mann, and Beech, 2003; Hanson, Broom, and Stephenson, 2004; Hanson, Steffy, and Gauthier, 1993; Harris, Rice, Quinsey, Lalumiere, Boer and Lang, 2003; Marques, Wiederanders, Day, Nelson, and van Ommeren, 2005; Quinsey, Khanna, and Malcolm, 1998; Rice, Quinsey, and Harris, 1991; Ruddiys and Timmerman, 2000; Schweitzer and Dwyer, 2003*).

One study (3%) provided qualified and partial evidence that treatment reduces recidivism (*Taylor, 2000*). One study (3%) found that treatment reduced violent recidivism but not sex offense recidivism specifically (*Lowden, Hetz, Harrison, Patrick, English, and Pasini-Hill, 2003*). Two studies (6%) suggest that changes in dynamic risk factors during treatment (and presumed to be the result of treatment) are correlated with reductions in recidivism (*Olver and Wong, 2009; Olver, Wong, Nicholaichuk and Gordon, 2007*).

Three studies (10%) did not offer statistical analyses of their findings (*West, et al., 2000 - data from "Kentucky", "New Hampshire", and "Vermont"*).

Three of the studies may be particularly notable because they are relatively recent, large-scale, and well-designed. The findings of what many regard as the best designed study of the efficacy of sex offender treatment, the Sex Offender Treatment and Evaluation Project in California, revealed no statistically significant differences in recidivism between offenders who were treated for 1-2 years and then released from treatment (with one year of parole supervision following treatment) and untreated sex offenders (*Marques, et al., 2005*).

The recent large scale study conducted by the Washington Institute for Public Policy (*Barnoski, 2006*) revealed that there was no reduction in sex offense recidivism for participants in the Washington Department of Corrections treatment program compared to those who did not participate in the program.

On the other hand, another recent and well designed study conducted in Minnesota (*Duwe and Goldman, 2009*) found that treatment produced "a significant albeit relatively modest reduction in sex offender recidivism".

The methodologies of many of the individual studies have been questioned making it even more difficult to reach definitive conclusions.

The conclusions of some of the individual studies suggesting a positive treatment effect are qualified by (sometimes unavoidable) design flaws. For example, two studies (*Looman, et al., 2000; Nicholaichuk, et al., 2000*) were criticized for having inadequate control groups (*Hanson and Nicholaichuk, 2000; Rice and Harris, 2003*). Two other studies (*Aytes, et al., 2001; Hildebrand, et al., 2004*) included as control groups sex offenders who had failed to complete treatment when it might have been equally appropriate to consider the failure of those offenders to be a poor outcome of a treatment endeavor rather than consider them as similar to offenders who never received treatment.

Another study showed a robust treatment effect (*Maletzky, et al., 2006*) but it should be noted that the treatment employed was hormonal/pharmacological, not the standard cognitive behavioral treatment employed by most sex offender treatment programs.

As already noted, one study which found no significant effect of treatment on sexual recidivism did find a significant positive effect of treatment on violent recidivism and overall recidivism (*Lowden, et al., 2003*). The authors concluded, in part, that sex crimes were the least commonly reported offenses making it difficult to compare groups using only measures of sexual recidivism.

It is worth noting that the authors of one recent analysis of studies concluded that the individual studies of treatment effectiveness that were more poorly designed or executed were more likely to show positive treatment effects than more well-designed and executed studies (*Hanson, Bourgon, Helmus and Hodgson, 2009*).

Some meta-analyses conclude that there is a small but significant effect of treatment on recidivism while some conclude no effect of treatment could be identified. The methodologies of many of the analyses have been subject to criticism.

Table 2 summarizes the findings of the meta-analyses surveyed




Table 2

Of the 8 meta-analyses I reviewed, 4 concluded that treatment had a small but significant positive effect on lowering recidivism (Gallagher, Wilson, Hirschfield, Coggeshall, and MacKenzie, 1999; Hall, 1995; Hanson, Gordon, Harris, Marques, Murphy, Quinsey, and Seto, 2002; Losel and Schmucker, 2005). One concluded that treatment reduced recidivism but that this was more true in weaker (poorly designed and executed) studies than in stronger ones (Hanson, et al., 2009). Two concluded that there was no such effect (Furby, Weinrott, and Blackshaw, 1989; Kenworthy, Adams, Bilby, Brooks-Gordon, and Fenton, 2004). One (Alexander, 1999) provided no statistical analysis.

Two meta-analyses (Gallagher, et al., 1999; Hall, 1995) have been critiqued about the particular studies which were chosen for inclusion. The Gallagher et al. meta-analysis has been criticized for including studies with “significant threats to validity” including early reports of studies which were contradicted by later versions of the same studies (Hanson, et al., 2004). The Hall meta-analysis has been criticized for including studies in which comparison groups were shown not to be equivalent; when these studies were removed from the analyses, the effect of treatment was no longer found to be significant (Hanson, Morton, and Harris, 2003; Harris, Rice, and Quinsey, 1998).

The authors of another meta-analysis (Losel and Schmucker, 2005) concluded that there is a significant effect of treatment on recidivism and that cognitive-behavioral treatments considered separately had a small but significant effect but much of the overall treatment effect appears to have come from studies in which treatment consisted of surgical castration.

Alexander's meta-analysis (1999) is sometimes cited as concluding that treatment is effective, probably because recidivism percentages derived from the analysis for treatment groups are somewhat lower than recidivism percentages for control groups, but, as noted, Alexander offered no statistical analysis of these findings.

The two meta-analyses in which it was reported that not significant treatment effect could be demonstrated have been criticized as well. The Furby, et al. analysis (1989) has been criticized because many of the studies included in the analysis focused on outdated treatment modalities. The Kenworthy, et al. analysis (2004) was withdrawn in 2008 pending an update.


Perhaps the most frequently cited recent meta-analysis is the ATSA Collaborative Outcome Data Project (Hanson, et al., 2002). The authors concluded that there was a small but statistically significant effect of treatment on sexual recidivism. It should be noted, however, that the difference in recidivism rates between treated and untreated offenders in the analysis was small (12% for treated and 17% for untreated). It should also be noted that the decision rules employed about study inclusion have been questioned and it has been suggested that the conclusions reached by the authors are not supported because many of the studies included in the meta-analysis had serious design flaws (Rice and Harris, 2003). Rice and Harris concluded that there is "no convincing evidence" that treatment is effective in reducing recidivism.

It should also be noted that some other reviewers, while not conducting a formal meta-analysis, have concluded that treatment is effective. For example, a review of studies conducted in 1991 by Marshall and colleagues concluded that there is "an unequivocally positive answer" to the question of whether sex offender treatment reduces recidivism (Marshall, Jones, Ward, Johnston, and Barbaree, 1991). However, others criticized this review saying that the authors' conclusion was not warranted because the review included too many studies that contained no adequate control groups and because the studies reviewed often could not ensure that comparison groups were equivalent (Quinsey, Harris, Rice, and Lalumiere, 1993).

In summary, research has not yet consistently demonstrated that time-limited treatment lowers sex offense recidivism in adults once they leave treatment and supervision programs.

Some individual research studies suggest that treatment lowers recidivism but others, including recent well-designed studies (and the only study to date employing randomized subject assignment), do not find such an effect. The findings of meta-analyses are sometimes encouraging but confusing and contested. Considered as a whole, it does not appear as of now that research persuasively demonstrates that time-limited treatment lowers sexual recidivism rates.

Is Treatment Effective?



Some approaches show promise.

A few studies suggest that recidivism rates may be lower for offenders who remain in treatment and supervision programs (“containment”) longer.

Some intervention or set of interventions may eventually be shown to lower recidivism in some offenders. For example, Marques et al. (2005) indicated that a combination of treatment response measures was found to be a significant predictor of sexual re-offense in high risk offenders suggesting that some form of treatment might eventually be effective with these offenders.

Prolonged treatment and intensive supervision (“containment”) may be important in lowering recidivism. The study of the Sex Offender Treatment Program at the Department of Corrections in Colorado revealed that re-arrest rates for all violent and sexual crimes (although, again, not specifically sexual crimes) were lower for offenders who remained in the DOC treatment program longer and who were subsequently placed on intensive parole supervision when released (Lowden, et al., 2003). It is interesting to note that Marques, et al. (2005) mentioned that such containment (English, 1998) might have improved the outcome of offenders in their study.

The study of the prison-based treatment program in Vermont revealed that both the length of time in treatment and the degree to which offenders received aftercare and correctional supervision following treatment contributed to lower recidivism (McGrath, et al., 2003). Another study found that providing offenders with increased support and opportunity to be accountable (through Circles of Support and Accountability Program) lowered recidivism following their release into the community (Wilson, et al., 2005).

The study of the community-based treatment program in Oregon indicated that the effect of treatment in lowering recidivism was particularly strong for offenders who remained in the treatment program for longer than a year (Aytes, et al, 2001).

A study conducted by the Colorado Department of Public Safety Division of Criminal Justice (2004) showed that high risk adult sex offenders living in a quasi-milieu setting which typically provides increased supervision and monitoring (“shared living arrangement”) had significantly fewer violations of probation than adult offenders with other living arrangements.

One meta-analysis indicates that adhering to “Need” and “Responsivity” principles may lower recidivism.

Hanson, et al., (2009) found in their meta-analysis that treatment programs which better targeted criminogenic needs and delivered treatment in a manner that was more likely to engage offenders were more effective in reducing recidivism.

July 2010

References

- Alexander, M. Sex offender treatment efficacy revisited. (1999). *Sexual Abuse: A Journal of Research and Treatment*, 11(2), 100-116.
- Aytes, K.E., Olsen, S.S., Zakrajsek, T., Murray, P., and Ireson, R. (2001). Cognitive/Behavioral treatment for sexual offenders: An examination of recidivism. *Sexual Abuse: A Journal of Research and Treatment*, 13(4), 223-232.
- Barnoski, R. (2006). Sex offender sentencing in Washington State: Does the prison treatment program reduce recidivism? Washington State Institute for Public Policy, Document No. 06-06-1205.
- Colorado Department of Public Safety Division of Criminal Justice (2004). *A report on safety issues raised by living arrangements for and location of sex offenders in the community*. Report dated March 15, 2004, retrieved from www.dcj.state.co.us/odvsom
- Davidson, P.R. (1984). *Behavioural treatment for incarcerated sex offenders: Post release outcome*. Paper presented at Sex Offender Assessment and Treatment Conference, Kingston, Ontario.
- English, K. (1998). The containment approach: An aggressive strategy for the community management of adult sex offenders. *Psychology, Public Policy and Law*, 4, 218-235.
- Duwe, G. and Goldman, R.A. (2009). The impact of prison-based treatment on sex offender recidivism. *Sexual Abuse: A Journal of Research and Treatment*, 21(3), 279-307.
- Friendship, C., Mann, R.E., and Beech, A.R. (2003). Evaluation of a national prison-based treatment program for sex offenders in England and Wales. *Journal of Interpersonal Violence*, 18, 744-759.
- Furby, L., Weinrott, M.R, and Blackshaw, L. (1989). Sex-offender recidivism: A review. *Psychological Bulletin*, 105, 3-30.
- Gallagher, C.A., Wilson, D.B., Hirschfield, P., Coggeshall, M.B., and MacKenzie, D.L. (1999). A quantitative review of the effects of sex offender treatment on sexual re-offending. *Corrections Management Quarterly*, 3, 19-29.
- Hall, G.C. (1995). Sexual offender recidivism revisited: A meta-analysis of recent treatment studies. *Journal of Consulting and Clinical Psychology*, 63, 802-809.
- Hanson, K.R., Bourgon, G., Helmus, L. and Hodgson, S. (2009). A meta-analysis of the effectiveness of treatment for sex offenders: Risk, need, and responsivity. Public Safety Canada, Publication 2009-01
- Hanson, K.R., Broom, I., and Stephenson, M. (2004). Evaluating community sex offender treatment programs: A 12-year follow-up of 724 offenders. *Canadian Journal of Behavioural Science*, 36, 87-96.
- Hanson, K.R., Gordon, A., Harris, A.J.R., Marques, J.K., Murphy, W., Quinsey, V.L. and Seto, M.C. (2002). First Report of the Collaborative Outcome Data Project on the Effectiveness of Psychological Treatment for Sex Offenders. *Sexual Abuse: A Journal of Research and Treatment*, 14, 169-194.
- Hanson, K.R., Morton, K.E., and Harris, A.J.R. (2003). Sex offender recidivism risk: What we know and what we need to know. *Annals of New York Academy of Science*, 989, 154-166.
- Hanson, K.R., and Nicholaichuk, T. (2000). A cautionary note regarding Nicholaichuk et al., 2000. *Sexual Abuse: A Journal of Research and Treatment*, 12, 289-293.

Hanson, K.R., Steffy, R.A., and Gauthier, R. (1993). Long term follow up of child molesters: Risk predictors and treatment outcome. *Journal of Consulting and Clinical Psychology, 61*, 646-652.

Harris, G.T., Rice, M.E., and Quinsey, V.L. (1998). Appraisal and management of risk in sexual aggressors: Implications for criminal justice policy. *Psychology, Public Policy, and Law, 4*, 73-115.

Harris, G.T., Rice, M.E., Quinsey, V.L., Lalumiere, M.L., Boer, D. and Lang, C. (2003). A multisite comparison of actuarial risk instruments for sex offenders. *Psychological Assessment, 15*(3), 413-425.

Hildebrand, M., deRuiter, C., and deVogel, V. (2004). Psychopathy and sexual deviance in treated rapist: Association with sexual and non-sexual recidivism. *Sexual Abuse: A Journal of Research and Treatment, 16*, 1-24.

Kenworthy, T., Adams, C.E., Bilby, C., Brooks-Gordon, B., and Fenton, M. (2004). Psychological interventions for those who have sexually offended or are at risk of offending. Cochrane Database of Systematic Reviews, Volume 3: CD004858. (Withdrawn in 2008).

Looman, J. Abracen, J., and Nicholaichuk, T.P. (2000). Recidivism among treated sexual offenders and matched controls: Data from the Regional Treatment Centre (Ontario). *Journal of Interpersonal Violence, 15*, 279-290.

Losel, F., and Schumucker, M. (2005). The effectiveness of treatment for sexual offenders: A comprehensive meta-analysis. *Journal of Experimental Criminology, 1*, 117-146.

Lowden, K., Hetz, N., Harrison, L. Patrick, D., English, K., and Pasini-Hill, D. (2003). *Evaluation of Colorado's prison therapeutic community for sex offenders*. A report of findings July 2003, Office of Research and Statistics, Colorado Division of Criminal Justice.

Maletzky, B.M., Tolan, A., and McFarland, B. (2006). The Oregon depo-Provera Program: A five-year follow-up. *Sex Abuse, 18*, 303-316.

Marshall, W.L. and Barbaree, H.E. (1988). The long-term evaluation of a behavioral treatment program for child molesters. *Behavior Research and Therapy, 26*, 499-511.

Marshall, W.L., Eccles, A., and Barbaree, H.E. (1991). The treatment of exhibitionists: A focus on sexual deviance, cognitive, and relationship features. *Behavior Research and Therapy, 29*, 129-135.

Marshall, W.L., Jones, R., Ward, T., Johnston, P., and Barbaree, H.E. (1991). Treatment outcome with sex offenders. *Psychology Review, 11*, 465-485.

Marques, J.K., Wiederanders, M, Day, D.M., Nelson, C., and van Ommeren, A. (2005). Effects of a relapse-prevention program on sexual recidivism: Final results from California's Sex Offender Treatment and Evaluation Project (SOTEP). *Sexual Abuse: A Journal of Research and Treatment, 17*, 79-107.

McGrath, R.J., Cumming, G., Livingston, J.A., and Hoke, S.E. (2003). Outcome of a treatment program for adult sex offenders. *Journal of Interpersonal Violence, 18*, 3-17.

McGrath, R.J., Hoke, S.E., and Vojtisek, J.E. (1998). Cognitive-Behavioral treatment of sex offenders: A treatment comparison and long-term follow-up study. *Criminal Justice and Behavior, 25*, 203-225.

Nicholaichuk, T., Gordon, A., Gu, D., and Wong, S. (2000). Outcome of an institutional sex offender treatment program: A comparison between treated and matched untreated controls. *Sexual Abuse: A Journal of Research and Treatment, 12*, 139-153.

Olver, M.E. and Wong, S.C.P. (2009). Therapeutic responses of psychopathic sexual offenders: Treatment, attrition, therapeutic change, and long-term recidivism. *Journal of Consulting and Clinical Psychology, 77* (2), 328-336.

Olver, M.E., Wong, S.C.P., Nicholaichuk, T. and Gordon, A. (2007). The validity and reliability of the Violence Risk Scale – Sexual Offender Version: Assessing Sex Offender Risk and Evaluating Therapeutic Change. *Psychological Assessment*, 19(3), 318-329.

Quinsey, V.L., Khanna, A., and Malcolm, P.B. (1998). A retrospective evaluation of the Regional Treatment Centre sex offender treatment program. *Journal of Interpersonal Violence*, 13, 621-644.

Quinsey, V.L., Harris, G.T., Rice, M.E., and Lalumiere, M.L. (1993). Assessing treatment efficacy in outcome studies of sex offenders. *Journal of Interpersonal Violence*, 8, 512-523.

Rice, M.E., and Harris, G.T. (2003). The size and sign of treatment effects in sex offender therapy. *Annals of the New York Academy of Science*, 989, 428-440.

Rice, M.E., Quinsey, V.L., and Harris, G.T. (1991). Sexual recidivism among child molesters released from maximum security psychiatric institutions. *Journal of Clinical and Consulting Psychology*, 59(3), 381-386.

Ruddijs, F. and Timmerman, H. (2000). The Stichting Ambulante Preventie Projecten method: A comparative study of recidivism in first offenders in a Dutch outpatient setting. *International Journal of Offender Therapy and Comparative Criminology*, 44(6), 725-739.

Schweitzer, R., and Dwyer, J. (2003). Sex crime recidivism: Evaluation of a sexual offender treatment program. *Journal of Interpersonal Violence*, 18, 1292-1310.

Taylor, R. (2000). A seven-year reconviction study of HMP Grendon therapeutic community. Home Office Research, Development and Statistics Directorate, Research Findings No. 115.

West, M., Hromas, C.S., Wengler, P., and Suthers, J. (2000). *State sex offender treatment programs: A 50-state survey*. Colorado Department of Corrections, August, 2000, retrieved from www.doc.state.co.us.

Wilson, R.J., Picheca, J.E. and Prinzo, M. (2005). Circles of Support and Accountability: An evaluation of the pilot project in south-central Ontario. Correctional Services of Canada, User report No. R-168.